

# 2003 Health and Preventive Practices Status Report

## from Missouri County-level Study



# Stone County\*\*

	County Prevalence (%)*	Southwest Region Prevalence (%)*	State Prevalence (%)*
<b>G</b> eneral			
Health Status (Fair or Poor)	19.0	18.8	16.9
No Health Coverage	21.0 H	17.4 H	12.3
<b>B</b> ehavioral Risk Factors			
Physical Inactivity	25.7	24.3	23.0
Current Smoker	34.4	26.8	26.5
Overweight (BMI 25-29.9)	41.8	35.6	35.3
Obese (BMI $\geq$ 30)	17.8	23.4	23.4
<b>D</b> isease and Conditions			
High Blood Pressure	20.1 L	27.9	28.5
High Cholesterol (35 years or older)	30.5	37.2	37.3
Asthma	11.3	11.6	10.7
Diabetes	6.1	6.4	7.2
Activity Limitation	13.3	18.0	17.8
<b>C</b> ancer Screening			
<b>Women</b>			
Never had a Mammogram	24.2	17.9	13.4
No Mammogram or Clinical Breast Exam in Last Year:			
40-49 years of age	68.1	52.3	49.4
50-64 years of age	52.1	43.3	35.9
65 and older	52.6	51.3	47.4
Never had a Pap Test:			
18-69 years of age	6.5	4.5	5.1
70 and older	0.0 L#	8.6	6.4
No Pap Test in Last Year (18-29)	45.2	28.1	25.8
No Pap Test in Last 3 Years (30-69)	7.1	13.8	13.8
<b>Men and Women 50 Years or Older</b>			
Blood Stool Test:			
Never had a Blood Stool Test	55.3	51.4	57.0
No Blood Stool Test in Last Year	73.9	74.2	77.5
Sigmoidoscopy/Colonoscopy			
Never had a Sigmoidoscopy or Colonoscopy	44.7	53.4	49.2
No Sigmoidoscopy or Colonoscopy in Past Five Years	59.7	63.3 H	57.0

\*Prevalence rates are age adjusted to the 2000 U.S. standard population

■ indicates that the prevalence rate is not significantly above or below state prevalence

H indicates that the prevalence rate is significantly above the state prevalence

L indicates that the prevalence rate is significantly below the state prevalence

Unless otherwise noted respondents are 18 years of age or older

\*\* The best indicators for Stone county were obtained by combining data from Stone and Taney counties.

# The number of respondents in this category is 0.